Math6502 test 11<sup>th</sup> January 2008

This test will last exactly 90 minutes.

Attempt all questions.

This is an open book test: lecture notes and text books are permitted in this exam. The use of an electronic calculator is permitted in this test.

- 1. (i) Write down the definition for the inverse of a matrix  $\underline{B}$ .
  - (*ii*) Consider the matrix:

$$\underline{\underline{C}} = \left( \begin{array}{rrr} 1 & 0 & -1 \\ 1 & 2 & 0 \\ 0 & 1 & x \end{array} \right).$$

For what values of x does the inverse of the matrix  $\underline{\underline{C}}$  not exist?

(*iii*) Consider the following system of linear equations:

$$2(x+y) = 1-z, x+z-2 = y, 2z = 5-y.$$

Write this system of equations in the form  $\underline{A} \mathbf{x} = \mathbf{b}$ , where  $\mathbf{x} = (x, y, z)$  and  $\underline{A}$  is a constant  $(3 \times 3)$ -matrix and  $\mathbf{b}$  is a constant three-dimensional vector.

- (*iv*) Either by inverting  $\underline{\underline{A}}$ , or using Gaussian elimination to express  $\underline{\underline{A}}$  in row echelon form [10] and using back substitution, find x, y and z and then verify your answer.
- 2. (i) Find the first three nonzero terms of the Maclaurin series for [5]

$$f(x) = \cos(nx).$$

(*ii*) Evaluate  $\frac{1}{\pi} \int_0^{\pi} x \cos(nx) \, \mathrm{d}x.$  [7]

## (*iii*) Sketch the periodic continuation, with period $2\pi$ , of the function [5]

$$f(x) = \begin{cases} -x & -\pi < x \le 0, \\ x & 0 < x \le \pi. \end{cases}$$

Is the resulting periodic function odd, even or neither?

(*iv*) Find the even Fourier series for f(x) with a periodic extension of  $2\pi$ . [8]

[5]

[2]

[8]

3. Let a one-dimensional iron bar of unit length, positioned between x = 0 and x = 1. The temperature of the bar is given by the function f(x, t). The initial temperature profile at time t = 0 is given by

$$f(x,0) = \begin{cases} x & 0 < x \le 1/2, \\ 0 & 1/2 < x \le 1. \end{cases}$$

Additionally the temperature of the bar is held at zero at both ends for all time.

- [2](i)Write down the one-dimensional heat equation in Cartesian coordinates.
- (ii)Using separation of variables, derive the general solution of the one-dimensional heat [10]equation to show that the general solution take the form

$$f(x,t) = C + Dx + \sum_{n=1}^{\infty} e^{-\kappa \lambda_n^2 t} \left( a_n \cos \lambda_n x + b_n \sin \lambda_n x \right) + \sum_{n=1}^{\infty} e^{\kappa \mu_n^2 t} \left( c_n e^{\mu_n x} + d_n e^{-\mu_n x} \right).$$

- (iii) Sketch the initial temperature profile.
- Neglecting unphysical solutions now let the general solution take the form (iv)

$$f(x,t) = C + Dx + \sum_{n=1}^{\infty} e^{-\kappa \lambda_n^2 t} \left( a_n \cos \lambda_n x + b_n \sin \lambda_n x \right).$$

By first applying the boundary condition at x = 0, then x = 1 show that the general solution has the form

$$f(x,t) = \sum_{n=1}^{\infty} b_n e^{-\kappa n^2 \pi^2 t} \sin n\pi x.$$

(v)By applying the initial condition solve the function f(x, t).

Let the matrices  $\underline{R}_{1}(\theta)$  and  $\underline{R}_{2}(\varphi)$  be given by

$$\underline{R_1} = \begin{pmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{pmatrix} \text{ and } \underline{R_2} = \begin{pmatrix} -\cos \varphi & -\sin \varphi \\ \sin \varphi & -\cos \varphi \end{pmatrix}.$$

- (i)What are the determinants of the two matrices?
- Using  $\cos(\alpha + \beta) = \cos\alpha\cos\beta \sin\alpha\sin\beta$  and  $\sin(\alpha + \beta) = \sin\alpha\cos\beta + \sin\beta\cos\alpha$ , [6](ii)express and simplify  $R_1 R_2$ . What is its determinant?
- Find  $\underline{R_1}^{-1}$  and  $\underline{R_2}^{-1}$  and hence or otherwise find  $(\underline{R_1}\underline{R_2})^{-1}$ . (iii)  $\left[5\right]$
- Are the two matrices,  $\underline{R_1}$  and  $\underline{R_2}$  symmetric? Show that  $\underline{R_1} \underline{R_2} \underline{R_2} \underline{R_1} = \mathbf{0}$ . (iv)[6]
- (v)What is the geometric interpretation of the two matrices  $\underline{R}_1(\theta)$  and  $\underline{R}_2(\varphi)$ ? [3]

4.

[5]

[5]

[5]

[3]